

Fecal Microbiota Transplantation

Policy ID:	HHO-DE-MP-1146
Approved By:	Highmark Health Options – Market Leadership
Provider Notice Date:	N/A
Original Effective Date:	N/A
Annual Approval Date:	01/2022
Last Revision Date:	10/27/2021
Products:	Medicaid
Application:	
Page Number(s):	1-3

Disclaimer

Highmark Health Options medical policy is intended to serve only as a general reference resource regarding coverage for the services described. This policy does not constitute medical advice and is not intended to govern or otherwise influence medical decisions.

POLICY STATEMENT

Fecal microbiota transplantation (FMT) involves the infusion of intestinal microorganisms via transfer of stool from a healthy person into a diseased individual, with the intent of restoring normal intestinal flora. Fecal transplant is proposed for the treatment of treatment-refractory *Clostridium difficile* infection (CDI).

DEFINITIONS

Highmark Health Options (HHO) – Managed care organization serving vulnerable populations that have complex needs and qualify for Medicaid. Highmark Health Options members include individuals and families with low income, expecting mothers, children, and people with disabilities. Members pay nothing to very little for their health coverage. Highmark Health Options currently services Delaware Medicaid: Delaware Healthy Children Program (DHCP) and Diamond State Health Plan Plus members.

POLICY POSITION

Fecal microbiota transplantation (FMT) may be considered medically necessary for treatment of individuals with recurrent CDI when BOTH of the following are met:

- There have been at least three (3) episodes of recurrent infection; and
- Episodes are refractory to appropriate antibiotic regimens, including at least one (1) regimen of pulsed vancomycin.

All other uses for FMT are considered experimental/investigational and, therefore, not covered because their safety and/or effectiveness cannot be established by review of the available published peer-reviewed literature.

ELIGIBLE PROCEDURE CODES

CPT Codes	Description
44705	Preparation of fecal microbiota for instillation including assessment of donor specimen.

44799	Unlisted procedure, intestine.
-------	--------------------------------

ELIGIBLE DIAGNOSIS CODES FOR PROCEDURE CODE 44705

Codes				
A04.71	A04.72			

References

Bafeta A, Yavchitz A, Riveros C, Batista R, Ravaud P. Methods and reporting studies assessing fecal microbiota transplantation: A systematic review. *Ann Intern Med.* 2017;167(1):34-39.

Cammarota G, Masucci L, Ianiro G, et al. Randomised clinical trial: faecal microbiota transplantation by colonoscopy vs. vancomycin for the treatment of recurrent *Clostridium difficile* infection. *Aliment Pharmacol Ther.* 2015;41:835-843.

Costello SP, Hughes PA, Waters O, et al. Effect of fecal microbiota transplantation on 8-week remission in patients with ulcerative colitis: A randomized clinical trial. *JAMA.* 2019;321(2):156- 164.

FDA. Important Safety Alert Regarding Use of Fecal Microbiota for Transplantation and Risk of Serious Adverse Reactions Due to Transmission of Multi-Drug Resistant Organisms 2019: <https://www.fda.gov/vaccines-blood-biologics/safety-availability-biologics/important-safety-alertregarding-use-fecal-microbiota-transplantation-and-risk-serious-adverse>. Accessed July 3, 2019.

Hayes, Inc. Emerging Technology Report. RBX2660 for Fecal Microbiota Transplantation. Lansdale, PA.: Hayes, Inc.; July, 2020.

Hayes, Inc. Emerging Technology Report. SER-109 for Fecal Microbiota Transplantation. Lansdale, PA.: Hayes, Inc.; July, 2020.

Hayes, Inc. Evidence Analysis Research Brief. Fecal Microbiota Transplantation for Treatment of Inflammatory Bowel Disease. Lansdale, PA.: Hayes, Inc.; July, 2020.

Hayes, Inc. Health Technology Brief. Fecal Microbiota transplant for Refractory or Recurrent *Clostridium Difficile* Infection in Adult. Lansdale, PA.: Hayes, Inc.; September, 2019.

Jiang ZD, Jenq RR, Ajami NJ, et al. Safety and preliminary efficacy of orally administered lyophilized fecal microbiota product compared with frozen product given by enema for recurrent *Clostridium difficile* infection: a randomized clinical trial. *PLoS One.* 2018;13(11):e0205064.

Kao D, Roach B, Silva M, et al. Effect of oral capsule vs colonoscopy-delivered fecal microbiota transplantation on recurrent *clostridium difficile* infection: A randomized clinical trial. *JAMA.* 2017;318(20):1985-1993.

McDonald LC, Gerding DN, Johnson S et al. Clinical practice guidelines for *clostridium difficile* infection in adults and children: 2017 Update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clin. Infect. Dis.* 2018;66(7).

National Institute for Health and Care Excellence (NICE). Faecal microbiota transplant for recurrent *Clostridium difficile* infection. London, UK: National Institute for Health and Care Excellence; 2014. NICE Technology Appraisal Guidance No. IPG484. Available at: www.nice.org.uk/guidance/ipg485. Accessed September 25, 2020.

Panchal P, Budree S, Scheeler A, et al. Scaling safe access to fecal microbiota transplantation: past, present, and future. *Curr Gastroenterol Rep*. 2018;20(4):14.

Quraishi MN, Widlak M, Bhala NA, et al. Systematic review with meta-analysis: the efficacy of faecal microbiota transplantation for the treatment of recurrent and refractory *Clostridium difficile* infection. *Aliment Pharmacol Ther*. 2017;46(5):479-93.

Staley C, Hamilton MJ, Vaughn BP, et al. Successful resolution of recurrent *Clostridium difficile* infection using freeze-dried, encapsulated fecal microbiota; pragmatic cohort study. *Am J Gastroenterol*. 2017;112(6):940.

Weingarden A, González A, Vázquez-Baeza Y, et al. Dynamic changes in short- and long-term bacterial composition following fecal microbiota transplantation for recurrent *Clostridium difficile* infection. *Microbiome*. 2015;3:10.

Youngster I, Russell GH, Pindar C, et al. Oral, capsulized, frozen fecal microbiota transplantation for relapsing *Clostridium difficile* infection. *Jama*. 2014;312(17):1772-8.

POLICY UPDATE HISTORY

<Date>	<Event>
--------	---------